



Entrant's Report

Harvest 2022 Oats YEN

YEN User ID: [REDACTED]

Entrant name: [REDACTED]

Main contact email: [REDACTED]

Sponsor/supporter: [REDACTED]

Sponsor/Supporter email: [REDACTED]

Field/Site name: [REDACTED]

Location: [REDACTED]


Incident energy 2019-20: 28 TJ/ha

Available water: 334 mm


Crop: Spring Oats

Variety: Husky






SUMMARY: A total of 11 Oats YEN entries were completed this season. Headline results for your entry are shown in benchmark diagrams below. Your yield of 7.3 t/ha ranked 6th within all Oats YEN entries. This represents 56% of its estimated yield potential of 13 t/ha, which ranked 5th within all Oats YEN entries in this season.



Overall yield rank:
6th



Overall potential yield rank:
5th



CONTENTS

Our detailed analysis of your yield result is provided in the following pages, including comparisons with other Oats YEN entries and with benchmarks taken from the Quaker Oats Growth Guide and the AHDB Nutrient Management Guide (RB209). We hope that this helps you to identify aspects of your husbandry and growing conditions that offer possible routes to further yield enhancement on your land.

Our approach in this report is to consider yield potentials and growing conditions for crops grown in this season, then the conditions of your crop, its development, its basic resources (light energy, water and nutrients), its success in capturing these and in converting them to grain. Lastly, we use grain analysis to provide a post-mortem on your crop's limiting components and nutrition.

CONTENTS	2
POTENTIAL GRAIN YIELDS	3
2022 Potential yields	3
SEASONAL GROWING CONDITIONS	4
In summary:	4
Soil description and nutrition analysis	6
Soil analysis	6
AGRONOMY	7
Variety	7
Husbandry factors	8
Husbandry factors continued	9
CROP DEVELOPMENT	10
RESOURCES AND THEIR CAPTURE	11
Water availability and capture	11
Energy capture	12
Nutrient capture	12
Image of this entry	16
YIELD ANALYSIS	17
Yield formation	17
Yield components	18
Grain formation and size	18
CROP NUTRITION POST-MORTEM	20
SUMMARY	22
The 2021-22 competition:	22
SPECIFIC COMMENTS ON THIS ENTRY	23
THE YIELD ENHANCEMENT NETWORK	24
CONTACTS	25
YEN SPONSORS	25

POTENTIAL GRAIN YIELDS



"The YEN exists to help you to enhance your yields."

The key to high yields amongst YEN entries has been called 'momentum' – maximising growth by avoiding setbacks. So, our approach to enhancing yields is to work out what limits growth – light energy, water, nutrients, or storage capacity – and then develop ideas to build better canopies, better roots, more storage, or better nutrition throughout growth.

To estimate potential yields, we assume a theoretically 'perfect' variety grown with 'inspired' husbandry on your land with this season's weather, achieving either:


- (i) **60% capture of light energy** through this season (including some in August), and its conversion to 1.2 tonnes of biomass per terajoule, or
- (ii) **Capture of all the available water** held in the soil to 1.5 m depth (or to rock if less) plus all rainfall from April to July, and conversion of each 18 mm into a tonne of biomass per hectare.

Taking the lesser of these two biomass amounts, we assume that a maximum of 55% can be used to form grain, this is the 'harvest index'. Note that we assume average temperatures for the UK, and no damage from waterlogging, frost, heat, or lodging.


The maps below show the potential grain yields for autumn and spring sown crops (i.e. before or after 1st Feb.) on retentive soils this season. For this we assume deep soils with no irrigation. They ranged from <12 t/ha upwards so, on most soils, high yields were theoretically possible almost everywhere.

2022 Potential yields

Autumn sown on retentive soil
(260 mm available water)



Spring sown on retentive soil
(260 mm available water)






We are using weather data from DTN™ this year. Note we do not have long term met data from DTN so cannot show a map of long-term average yield potentials.

SEASONAL GROWING CONDITIONS

In summary:


The adjacent graphs show the monthly temperatures, rainfall and total solar radiation for your area through this growing season compared to your regional long-term average (LTA) and the average for all UK arable areas (1981-2010, from the Met Office).

- Autumn 2021 allowed timely drilling; then a warm winter enabled good establishment.
- Residual herbicides were effective, but weeds emerged and grew through the winter so post emergence herbicides were needed. Groundsel was an issue.
- Slug pressure was strong, especially in the South East.
- The year was warm throughout, particularly May to July in the East so crops were advanced throughout and spray times were 7-14 days earlier than normal.
- May saw less sunshine than normal but June and July were bright and hot, so overall solar radiation for the year was 7% above the 5 year average.
- Summer was drier than in any other YEN year (since 2013) with 33% less rain normal so most crops became water-limited through June and July, particularly on light or shallow soils. This and the heatwave, with record temperatures in early July, caused early senescence of many crops and an early harvest.
- Winter oats diseases: Crown rust pressure began to build in early June, although there was some regional variation. High pressure was recorded in untreated trial plots in the south-west, although commercial fungicide programmes provided effective control for most growers. Powdery mildew was typically contained to the base of the canopy.
- Spring oats diseases: The dry spring weather limited the development of foliar disease, with low levels of crown rust and powdery mildew recorded around the time of T2 applications. These infections were well controlled by most fungicide programmes and crops remained relatively clean throughout the season.



YEN Benchmarking charts – What do they mean?

YEN is much more than a competition – it provides a full set of metrics whereby you can gauge the performance of your crop against all other YEN crops. This has provided the principle value of YEN to most participants. We do this with benchmark charts. These compare your value with everyone else's entered in the Oats YEN this year and with standard benchmarks, if available and appropriate. The key is as follows:



The 'whiskers' show the range of Oats YEN values this season whilst the grey box shows the middle half of values, with a line for the mid-value. The orange line shows the value for your entry. Blue and green lines indicate benchmark values for winter and spring oats, respectively, from the Quaker Oat Growth Guide. Benchmark charts throughout this report summarise data provided for all YEN Oat crops (they exclude barley, wheat, rye and spring wheats).

Note that 'Dynamic Benchmarking' is available to all YEN members via the YEN website. This means you can compare your own yield or grain nutrient data with subsets of all other YEN crops selected by crop type, soil type, location or year back to 2013.

As Oats YEN entry numbers are relatively low, all data from both winter and spring Oats entries have been combined into one box and whisker plot per parameter. This may mean a wide range of values are apparent for some parameters. As the competition grows, it may be possible to display separate figures for winter and spring Oats, making them more applicable to your entry.

Soil description and nutrition analysis




Your soil's capacity to hold available water is critical in determining your potential yields. We rely on entrants describing the soil where their YEN entry grew. We can use the [UK Soil Observatory map viewer](#) to check whether this complies with the surrounding land.

Good soil descriptions are vital in allowing us to estimate soil water holding capacity and, along with summer rainfall, the water available to your crop (see Benchmark charts in the section on 'Resources & their Capture').

Topsoil analyses provided by NRM also tell us about soil status for pH, P, K and Mg, as reported on the next page. A few sites show low values for soil pH, P, K or Mg. If these are unexpected, they may need further checks, either by repeating soil analysis and by checking both leaf and seed analyses later in this report. Previous YEN leaf and seed nutrient data have indicated that UK cereal crops often experience deficiencies in one or more nutrients, and sometimes this is despite soil levels being satisfactory. So, by combined use of soil, leaf and seed analysis, the YENs now help to diagnose whether nutrient shortfalls are arising from poor supply, or poor capture by the root system.

Soil analysis



Soil pH <6 is acid. High pH soils may require that special attention is paid to phosphorus (P) and micro-nutrient levels in leaf and grain (see later).

Only a small difference separates P Index 0 (≤ 9) and 2 (≥ 16). High yields are possible at P index 1 but fresh P is also usually required. Use grain P (see page 20) to check if P was sufficient.

Soil potassium (K) analysis checks on whether K supplies are likely to have been deficient for average crops. However, high yielding crops require very large amounts of K.

Magnesium (Mg) is a key component of chlorophyll so deficient plants show striking inter-veinal yellowing. Temporary deficiencies often occur in springs if topsoils are dry.

SOM supports crop performance through better nutrient availability, soil aggregation, and water holding capacity. NRM determines SOM by 'loss on ignition'. Note: other methods can give lower values.


AGRONOMY

This section considers how your variety and husbandry decisions related to others entering the YEN this year. Note that the multi-year YEN dataset suggests that the individual effects on grain yield of variety choice or husbandry decision are relatively small; it is how these decisions (and other factors) are combined into the overall strategy on each farm that is responsible for the level of yield that tends to be achieved. Hence it should be possible to learn from the best performing farms. In summary, we are concluding that:

- **High yields** are not restricted to just one part of the UK.
- **Attention to detail** is important.
- **Large yields come from large crops** . . . i.e. taller crops with more fertile shoots
- **Best yielding seasons** had dry, bright autumns and winters, bright springs and cool summers
- **Good nutrition is hard:** most crops suffer nutrient deficiencies, especially of P.





Variety

Oats YEN entries in this season included 5 different Oats varieties. Your variety was Husky, which according to the AHDB Recommended List (or alternative source for some varieties) 0 and has an average specific weight of kg/HI




Husbandry factors

The following diagrams use orange segments or orange bars to indicate the agronomy of your crop, if known, so you can see how this relates to all other YEN entries.




Husbandry factors continued


Sowing date: Winter




Seeds sown per m²




Sowing date: Spring




Number of PGRs applied




Total N applied, kg/ha




Number of herbicides applied



Number of N applications




Number of insecticides applied




Fertiliser P₂O₅ applied, kg/ha




Number of fungicides applied




Fertiliser K₂O applied, kg/ha




Crop protection spend, £/ha



Fertiliser SO₃ applied, kg/ha



Crop protection spend, £/tonne




CROP DEVELOPMENT

The following charts show how your entry developed through the season, compared to all other winter and spring oats YEN entries and Benchmarks. The cardinal stages of emergence (GS10), start of stem extension (GS31), flowering (GS61) and full senescence (GS87) determine the length of each phase for growth:


- Foundation, GS10-GS31 – when tillers and main root axes are formed,
- Construction, GS31-GS61 – when yield-forming leaves, ears and stems are formed, including soluble stem reserves
- Production, GS61-GS87 – when grains are filled, both with new assimilates and reserves redistributed from stems.

Emergence date




The wide range of emergence dates is due to the inclusion of winter and spring oats crops. Full emergence in oats is normally completed >300 day degrees (oC days) after sowing. Deep sowings take longer.

Stem extension (GS31)




Stem extension triggers faster growth because the stem provides a new sink for assimilates.

Flowering (GS61)




Flowering signals the start of grain formation. Delays in flowering, due say to cold weather after ear emergence, may cause growth to pause.

Canopy senescence (GS87)




No further growth can occur after the canopy has fully senesced.

Harvest date



Harvest dates are highly susceptible to rain patterns through August and September.

Crop height, cm




We measure height on the harvest 'grab' samples. We omit measuring samples which look to have been cut above ground level.

RESOURCES AND THEIR CAPTURE

Water availability and capture


This page shows how weather this year affected the water available for your crop and other crops entered in the Oats YEN. Water is supplied through the main growing period from concurrent rainfall and also from water stored in the soil. UK soils almost always refill with water over-winter. Water potentially available to each crop through the summer includes all this soil water plus the summer rainfall (April to July).

Soil water holding capacity, mm



Deep soils hold water to a great depth; we assume roots can access all easily held water (to 2 bar suction) to a depth of 1.5m (or to rock, if shallower). If enough roots didn't reach to this depth, capture of soil-available water will have been accordingly less.


Rainfall April–July, mm



After winter drainage stops, spring and summer rainfall is held in the topsoil until it is evaporated or transpired by the crop's canopy.


Whilst we cannot yet measure water captured by YEN crops individually, by assuming your crop's conversion of water to total biomass was 'normal' (20 mm water for each t/ha biomass formed), we have made crude estimates below of the likely success of your crop's root system in capturing water.

Total water available, mm



Total water is the sum of your soil's water-holding capacity and your summer rainfall (both shown above).

Estimated use of available water, %



Small water use will sometimes have been due to less demand for canopy transpiration (e.g. because crop developed faster and matured earlier) or otherwise due to worse rooting.


If your estimated use of available water exceeds the total water available, this may be good news! It either suggests that your crop's roots were more efficient than normal, or that your soil description was overly pessimistic: i.e. your soil apparently managed to provide more water than we estimated was possible from your soil's texture, stone content and depth.

A high yielding crop, growing say 20 t/ha of biomass (so yielding 12 t/ha grain at 51% harvest index), would need to capture ~400 mm from soil plus summer rain. Given that most of the UK's arable area often receives only 200-250 mm summer rainfall (from April to July), a large proportion of the water for high yielding crops must come from that held in the soil since the winter, mainly in the subsoil.


Energy capture

The benchmarking charts below show how weather this year affected light energy available for this entry and other Oats YEN crops.


Solar radiation Oct–Mar, TJ/ha



Solar radiation Apr–May, TJ/ha




Solar radiation Jun–July, TJ/ha



Solar radiation in September last year and August this year has been omitted, because few crops were green during those months, but crops could have achieved greater total biomass, and possibly also grain biomass, if they maintained green canopies during any part of these two months.


Whilst we cannot yet measure light capture by YEN crops individually, by assuming your crop's conversion of light-energy was 'normal' (1.2 tonnes/TJ), we have made a crude estimate below of the likely success of your crop's canopy in capturing total light-energy for the 12 months of this season.

Solar radiation total, TJ/ha/yr



Total solar radiation across YEN entries is generally less in the north and more in the south.

Estimated % solar radiation captured











Average light capture tends to be poor if a crop's lifespan is short. The benchmark winter and spring oats crops intercept 55% and 49% of available solar radiation, respectively.

Nutrient capture

Whether nutrient capture was sufficient to support full conversion of light and water is best deduced from nutrient concentrations in crop tissues – both leaves (next three pages) and grains (later section). No critical thresholds or benchmarks are shown for leaf analyses because these change through a crop's life and are still uncertain. However, the benchmarking diagrams should enable you to compare your crop's levels with all other YEN entries this year, analysed at the same time.

Lancrop Laboratories provide leaf analyses for YEN. Samples are of the newest fully expanded leaf.











Image of this entry

Images are a very efficient way of collecting lots of information. An overhead photo taken during grain filling gives an impression of canopy size, nutrition and health.




YIELD ANALYSIS

Yield formation


The whole-crop samples that YEN entrants provide all have their components counted and weighed and results are shown in the following charts, assuming that each sample was representative of the whole area from which grain yield was determined. [All area-related values are derived from the validated grain yield.]

Total biomass production indicates the success with which a crop captured its key resources, light-energy and water, and the harvest index (the proportion of total biomass that was harvestable) indicates how this biomass was apportioned to grain. Since grain growth happens last, harvest index also indicates how late growth related to early growth.


Your grain yield (expressed as t/ha and % of potential) is shown below along with biomass and harvest index, in relation to all other Oats YEN entries and to the Oat Growth Guide Benchmark yields of 8.0 t/ha for winter oats and 7.0 t/ha for spring oats.




YEN experience has been that high biomass relates to high yields.




Harvest Index is the percentage of total biomass that was harvestable as grain. Years with high fertile shoots tend to have low harvest index.



Benchmark yields of winter and spring oats are 8.1 and 7.0 t/ha, respectively. Northern crops tend to yield slightly higher than those grown in the south.



YEN yield potential reflects light energy and water available at your site this year, expressed in t/ha.




Any YEN entry exceeding 100% of its estimated potential must have found more light or water than was estimated at this site, or must have grown with exceptional efficiency.

Yield components


Grab sample analysis tells the story of your crop because the different yield components are determined sequentially, first shoots, then panicles, then grains per panicle, then grain weight. Comparing your yield components with those of other YEN entrants should indicate the stage(s) through the season at which your crop deviated from others and from normal (represented by the AHDB Benchmarks, blue lines).

Panicles No./m²




Panicles per m² depend on plant establishment, then tillering, and then the survival of each shoot during stem extension.

Spikelets per panicle




Spikelet numbers are determined between GS30 (panicle at 1 cm) and GS31 (1st node). Numbers are crucial for barley but not oats because oats spikelets are flexible in the number of grains they set, so they can compensate.

Grains per panicle




Grains per panicle can be less than normal either through compensation for high ear numbers or if May was dull.

Grains '000s/m²



High yields almost always depend on grain numbers per m² being high through combining good panicle numbers with adequate grains per panicle (above).

Grains set per g chaff




High grains/g chaff indicates that conditions around flowering time were good for photosynthesis.

Grain formation and size


We use your combine-harvested grain sample to provide the analysis of grain size and grain filling on the next page. Grain filling depends mainly on photosynthesis after flowering, therefore it largely relies on the health and longevity of the green canopy, but sugars stored in the stem can also provide 2-4 t/ha of assimilates for grain growth and most of the protein from senescing leaves is also redistributed to form grain protein.

We have not measured stem sugars in YEN so far, but it is possible to assess them using a refractometer (giving a so-called Brix reading). Stem storage of sugars depends on shoot numbers and sunshine levels in May being good.


If grain number per m² is low (see above), or if conditions during early grain-fill are limiting, final grain filling, hence yield, may be constrained even if later conditions are good – this is sometimes described as 'sink' limitation. We try to use analysis of grain volume and grain density to deduce whether crops were sink limited.




Thousand grain weights (TGW) depend on early grain expansion to set up the potential grain size and then on continuing supplies of photosynthates to replace grain water with starch.




Specific weight is an important quality parameter for oats.




Grain length is set before grain width and tends to indicate potential grain storage capacity.




Grain width reflects the success with which grain storage capacity is filled.




A high grain L:W ratio indicates that assimilate supply for grain filling did not fully satisfy the grains' potential storage capacity.



Grain volume here is the product of length and cross-sectional area, assuming grains are ovoid.



Low density grains indicate that grain filling was not constrained by storage capacity (volume) – so they were not 'sink limited'.





Did you know more than half of a load of grain is air?! High specific weight is achieved by having both dense grains and small voids between grains (under standard packing conditions).

CROP NUTRITION POST-MORTEM


The YEN has trail-blazed use of grain analysis to provide overall post-mortems on a crop's nutrition.

- Results from >900 YEN cereal samples analysed up to last year suggest that nutrient deficiencies are very common (using the 8 critical values that we know so far); >80% of crops showed deficiencies, and >50% showed two or more deficiencies! Phosphorus deficiency has been most common.
- YEN Nutrition was therefore launched in 2020 to help to remedy these deficiencies – further details and registration are available [here](#)
- Crop nutrients differ in how they are shared between grain and straw at harvest. The graph below shows how different crop species store most of their N & P in the grain but most of their K in the straw (as estimated from analyses of feed materials).



A wheat grain showing where each of the 12 essential nutrients is most concentrated.

- We now use YEN-low values (i.e. lower quartiles from all past YEN data – the boundary between the bottom quarter and top three-quarters of all YEN values since 2013) as comparators (red lines) for all nutrients in all crops. We find YEN-low values to be very similar to known critical thresholds of N, P, S and Mn in wheat, as well as to less certain critical values of K, Mg, S & Zn, so we assume they can be applied for all nutrients in all crops.
- The following benchmarking-charts and YEN-low values provide the best means of identifying the nutrient(s) most likely to have limited your crop.
- Critical grain protein (or N% x 5.7) levels are variety-dependent so it's best to compare your value with the value reported in the AHDB Recommended List for that variety. If the observed protein level is significantly less or more than the RL value, we take this to indicate that this crop was under- or over-supplied with nitrogen.



In oats, grain protein (N% x 5.7) is less often used as a specification for grain marketing than other cereals, but is an indicator of the success of the nutrition strategy. Benchmarks are 11.6% and 14.8% for winter and spring oats, respectively.

Recent research has shown grain P analysis can provide a useful check on sufficiency of phosphorus. Many past YEN values have been less than 0.32% indicating the difficulty of ensuring good P supply and capture.

RB209 assumes a standard value of 0.54% potassium (K) in grain. Values less than 0.38% indicate a need for further checks on K nutrition, especially by soil analysis but also by analysing leaves.

Calcium nutrition relates to the crop's use of water. However, almost all the crop's calcium remains in the straw at harvest, so we are yet to learn whether grain calcium can tell us about the crop's water status.

Literature shows low magnesium (Mg) values in grain are <0.08%. YEN data from previous years show high grain Mg has been associated with high grain yields.

S is required in proportion to grain protein formation.

A high N:S ratio (greater than about 17) indicates the crop was affected by sulphur deficiency.

Literature shows low manganese (Mn) values in grain are <20 mg/kg. If crops show low grain manganese, leaf Mn should be checked.

Grain copper (Cu) less than 2 mg/kg indicates possible deficiency. Very few YEN samples have been this low.

Zinc (Zn) deficiency has been rare in YEN crops. Values below 15 mg/kg are considered deficient. Grain zinc appears to inter-relate with nitrogen availability.

We currently have no guidelines for grain iron (Fe) interpretation. Average Fe has been around 40 mg/kg in previous years of YEN.

Most Boron is kept in the straw at harvest. Previous YEN boron values have varied hugely with season, so grain analysis may not be useful for assessing boron sufficiency.

SUMMARY

The 2021-22 competition:

- Many congratulations for providing the information necessary to complete this report; the collective efforts of all YEN contributors serve to maximise the value of what can be reported and the deductions that can be made for everyone – we call this approach 'learning by sharing' and believe that the whole industry would benefit by making this approach their normal practice.
- The number of participants in Cereal YEN this year was more than last year, and similarly diverse, so we have been able to report and rank barley and oats entries separately. Crops were from all over the UK, and from across Europe including the Netherlands and Denmark. All these data are invaluable; the more data we have, the more robust and confident we can be in the comparisons we make, both when 'benchmarking' and when analysing associations within the whole set of data.
- This was the 10th year of YENs and the 2nd year of the Oats YEN. The winning field yield in 2022 was a new YEN record of 18.0 t/ha (in Lincolnshire). As each YEN year passes and as more YENs develop, we are increasingly struck by the farm-to-farm differences; some farms are consistently achieving high yields, and several farms have achieved YEN Awards over several seasons. It is evident that a 'farm factor' is playing a big part in governing yield levels. This gives real value to being a YEN participant – through having an opportunity to compare with and learn from others who consistently perform very well.
- Estimated UK farm average yields in 2022 were 7-9% better than the 5-year average for each of the cereals, but only wheat (and possibly oats – we only have data since 2020) showed better YEN yields this year. YEN yields far exceeded farm averages:

Cereal yields in 2022	Winter wheat	Winter Barley	Spring Barley	Oats
Defra farm yield estimate, t/ha	8.59	7.42	6.22	5.79
Change from previous 5 years	+8%	+7%	+9%	+8%
Average YEN yield in 2022, t/ha	11.31	9.20	7.38	6.95
Change from previous 5 years	+6%	-1%	-3%	(+8%)

- In terms of physiology, over all ten years of the YEN high yields associate with high ear numbers and high total biomass; the latter is more important than harvest index in explaining yields. Hence the importance of striving for good light and water capture.
- In 2022, after good crop establishment and tillering, the dry spring restricted N uptake so shoot survival was reduced. Then a dull May did not allow floret survival to compensate, so grain numbers were less than the YEN averages. However, flowering and grain filling were about a fortnight ahead of normal and crops were surprisingly successful in capturing soil-stored water. Hence, grain filling largely escaped the worst of the hot, dry July and more than compensated for the reduced grain numbers. Thus, YEN yields were slightly better than the long term average.
- In summary, yields in 2022 were good. Because of their generally widespread moisture limitation yields showed many growers just how much water their soils and rooting systems can provide. This should be very useful in building yield enhancing strategies into the future.

Comments on the next page are generated automatically from your data, with the aim of high-lighting features of your crop which may point out routes to yield-enhancement on your land.

SPECIFIC COMMENTS ON THIS ENTRY

Resource capture, growth and yield:

- High YEN yields have generally been associated with high biomass production. Your yield arose from a low total biomass and a normal harvest index.
- Our target for annual light interception by annual crops (whether sown in autumn or spring) is 60% compared with 36.4% achieved by this crop.

Crop Nutrition:

- Your crop took up 125 kg N/ha. The benchmark N uptake at harvest is 205 kg N/ha for winter oats and 157 kg N/ha for spring oats.

Review of Oilseed YEN

Out of 45 entries in 2022 gross output of oilseed rape crops ranged from 3.4 t/ha to 8.9 t/ha, with the winning crop located in Lincolnshire achieving an impressive 8t/ha yield along with an oil content of 48.5%. On average crops achieved 61% of potential yield which ranged from 7.0 t/ha to 12.5 t/ha. Crops generally established well, with adequate moisture minimising CSFB damage. Dry conditions in spring would have restricted canopies becoming too large, although conditions may have hampered N uptake in some crops. Sunny April conditions along with rainfall in May were conducive for setting high seed numbers. The majority of seed filling took place in June which was sunnier and only slightly warmer than average - giving a high rate of photosynthesis helping seed filling and oil formation. Oilseed YEN also included 10 Linseed entries this year – it has been great to see the number of entries increase and develop our understanding of linseed physiology and yield potential.



Wheat Quality Competition

The YEN Wheat Quality Award, sponsored by UK Flour Millers, is taking place again in 2022. All Group 1 wheat entries which provided a large grain sample have been entered and the best are being short-listed. Following breadmaking analysis and assessment, the winners will be announced during the Awards Dinner evening on Tuesday 24th January 2023 (see below).



AHDB events

Several AHDB Monitor Farms entered the YEN competition for 2022 and YEN will be included in a number of upcoming monitor farm meetings, please visit the [AHDB website](#) for more details.



YEN Nutrition

YEN Nutrition was initiated in 2020 because YEN data had indicated that the majority (>80%) of crops have inadequate nutrition, one way or another. This new YEN connects anyone – farmers, advisors, suppliers and academics in the UK or abroad – seeking to improve nutrition of any grain crop – cereal, oilseed or pulse. Membership begins with grain analysis and grain nutrient benchmarking on six or more fields. Further details are available [here](#).



YEN Zero

With the industry targeting net zero emissions by 2040, YEN Zero was initiated in 2021 with the overarching aim of creating a community of farmers and key players from across the agricultural industry to compare field-scale carbon footprints across farms, and build understanding of the issues and opportunities for making progress. Further details are available [here](#).



'YEN is ten' Conference and Awards dinner

If you haven't done so already, please register and come to the YEN's tenth birthday conference at East of England Arena, Peterborough on 24th January 2023. You can Register for the YEN Conference [here](#).




CONTACTS

Please send any comments, observations or queries to the contacts below.

Dhaval Patel	Dhaval.Patel@adas.co.uk	07502 658098
Sarah Kendall	Sarah.Kendall@adas.co.uk	07720 496793
Roger Sylvester-Bradley	Roger.Sylvester-Bradley@adas.co.uk	07884 114311
Daniel Kindred	Daniel.Kindred@adas.co.uk	07774 701619
Sarah Clarke	Sarah.Clarke@adas.co.uk	07890 532581

Or email yen@adas.co.uk for general enquiries.

 @adasYEN

YEN SPONSORS

The YEN was initiated by industry and is entirely industry funded. We are most grateful to all our sponsors. They not only provide funding but they are fundamentally involved in management of the YEN and in supporting individual farms in making their YEN entries. The YEN would not exist without them!



Visit www.yen.adas.co.uk for sponsors' details, news updates and to register for 2021.